



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : A01N 1/02, 25/34	<b>A1</b>	(11) International Publication Number: <b>WO 94/13135</b> (43) International Publication Date: 23 June 1994 (23.06.94)
(21) International Application Number: PCT/EP93/03364 (22) International Filing Date: 1 December 1993 (01.12.93) (30) Priority Data: MI92A002776          4 December 1992 (04.12.92)          IT (71) Applicants (for all designated States except US): DEVEL- OPMENT BIOTECHNOLOGICAL PROCESSES S.N.C. DI PELLICCIA MARIA TERESA [IT/IT]; Via Beata Francesca, 10, I-83100 Avellino (IT). CONSORZIO PER LA GESTIONE DEL CENTRO DI BIOTECNOLOGIA AVANZATA [IT/IT]; Viale Benedetto XV, 10, I-16132 Genova (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): GERACI, Giuseppe [IT/IT]; Via Botteghe, 162, I-80046 S. Giorgio a Cremano (IT). DE ROSA, Mario [IT/IT]; Via E. Nicolardi, 188, I-80131 Napoli (IT). ROSSI, Mosè [IT/IT]; Via E Nicolardi, 188, I-80131 Napoli (IT). CANCEDDA, Ranieri [IT/IT]; Viale Benedetto XV, 10, I-16132 Genova (IT). DE LUCA, Michele [IT/IT]; Viale Benedetto XV, 10, I-16132 Genova (IT). PELLEGRINI, Graziella [IT/IT]; Viale Benedetto XV, 10, I-16132 Genova (IT).		(74) Agent: MINOJA, Fabrizio; Studio Consulenza Brevettuale, Via Rossini, 8, I-20122 Milano (IT). (81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: CRYOPROTECTIVE AQUEOUS SOLUTIONS USEFUL FOR THE PRESERVATION OF <i>IN VITRO</i> CULTURED EPITHE- LIAL SHEETS (57) Abstract <p>A method for the preservation of <i>in vitro</i> expanded epithelial sheets which comprises the incubation of said sheets in a cryopreserving solution characterized by lack of volume increase at the liquid-solid phase transition. The cryopreserving solutions comprise: a) polyethylene glycols having molecular weight not higher than 20 KD; b) at least one cross-linking agent selected from polyols, polyamines, mono- or oligosaccharides, polyethyleneglycols having molecular weight lower than 1 KD.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

CRYOPROTECTIVE AQUEOUS SOLUTIONS USEFUL FOR THE  
PRESERVATION OF IN VITRO CULTURED EPITHELIAL SHEETS

The present invention refers to the preservation, by freezing, of epithelium sheets obtained from cell cultures of keratinocytes.

The cultures of human keratinocytes, developed by  
5 Green and Rheinwald (US 4,016,036; US 4,304,865; US  
4,456,687) raise more and more interest in several  
medical applications such as in the treatment of  
serious burns, ulcers, difficult to heal scars and in  
plastic oncological and reconstructive surgery.

10 In the case of seriously burnt patients, the  
death-rate, depending on the patient's age and affected  
body surface, is dramatically high, particularly for  
third degree burns greater than 60%. According to the  
usual procedure, the damaged body surfaces are covered  
15 by transplanting skin films removed from healthy areas  
(auto-grafts); this procedure is very time-consuming  
and cannot be practically used when the burnt area  
exceeds 60% of the body surface. The technique by  
Rheinwald and Green has allowed the burnt areas to be  
20 covered with autologous epithelial films obtained from  
the patient's epithelial cells (keratinocytes).  
Starting from a small biopsy of epithelial tissue, the  
technique affords an epithelium surface sufficient to  
cover the whole body surface in a few weeks.

25 The autologous transplant in seriously burnt  
patients represents a unique, socially relevant  
therapeutic approach.

Autologous transplants are carried out also in cases of plastic oncological and reconstructive surgery. In this case, before surgery, sheets of epithelium expanded in vitro are prepared starting from a patient's biopsy.

The use of cultured human epithelium sheets as "living bandage", for the treatment of ulcers and scars which are difficult to heal, is particularly interesting not only from the social but also from the commercial point of view. In this case, the epidermis sheet, optionally obtained from suitably typized keratinocytes cultures, is used as a temporary dressing of the scar, making the restoring of the epithelium thereon easier thanks to the production of a series of factors stimulating the growth of this tissue, decreasing therefore the patient's suffering and hospitalization.

More than 4 million patients could be treated in the US with this new therapeutic approach. It is obvious however that this would ask for industrial premises, remarkably reduced costs of the epithelium sheets, long and easy preservability of the product and easy application procedures in hospital environments.

The method originally developed by Green and Rheinwald for autografts comprises the sampling of a small fragment (2-3 cm<sup>2</sup>) of epithelium from a healthy area, from which an epithelial cellular suspension is obtained and cultured and finally expanded by means of subsequent passages of the primary cultures. When the secondary and tertiary cultures become confluent,

multilayered epithelium sheets are obtained in different sizes according to the need.

Notwithstanding the high potentiality of this therapeutic approach, its wide-scale application is limited by the need that the obtained epithelial sheets should be used within a few hours after their preparation.

EP-A-0296475 discloses a method for preserving at  $-100^{\circ}\text{C}$  viable, transplantable epithelium sheets cultured in vitro, frozen in a nutrient medium containing the usual cryopreserving agents such as dimethylsulfoxide or glycerine.

Although this method has allowed for the first time the preservation of in vitro cultured epithelium sheets, a number of problems limit its industrial applicability:

- a) the permanence in the preserving medium before freezing is extremely critical for the cells at room temperature. This makes the initial step of the preserving process, comprising an incubation at room temperature to equilibrate the epithelium sheets with the solution and the final phase of thawing before using the sheets, critical. In both cases, a few minutes difference in the performance of these steps may (irreparably) impair the characteristics of the product;
- b) The complexity of the cooling cycle of the samples, critically characterized by different cooling rates in particular temperature ranges involve expensive industrial apparatuses and high

production costs;

- c) The preserved product is very sensitive to even slight changes in the preservation temperature, which can cause alteration of the product preventing its use;
- d) The shipment of the sample to the utilization sites must necessarily be carried out at a temperature constantly kept at about  $-80^{\circ}\text{C}$  and similar measures must be observed by the hospitals before the use of the product;
- e) Cell viability in the frozen graft obtained by this procedure can be in the best conditions up to 70%. However given the critical steps required for freezing (see points A, B, C, D) the large variability of the obtained results makes this procedure difficult to use in large scale preparations.

The present invention aims at overcoming the above limits, so as to allow the industrial applicability of the production and preservation processes of human epithelium sheets optionally typized, convenient methods of use in hospital environments and costs compatible with the previously cited therapeutic uses.

The above problem is solved by the present invention which provides a new kind of cryoprotective solutions allowing to preserve for a long time frozen cultured human epithelial sheets.

According to the invention, the processing times are not critical both before freezing and in hospitals where thawing and grafting take place.

Moreover, the quality of graft frozen with the new procedure is much superior cell viability can be up to 100% in the best conditions. At variance with grafts frozen with the old procedure the frozen sheets, preserved in these new cryoprotective solutions, do not undergo functional alterations impairing their use, even if the preservation temperature is subjected to changes of tens of degrees or the sample is accidentally thawed for a short period of time.

It is known that in the cryopreservation of biological material in aqueous solutions, the formation of ice crystals, connected with the regular growth of the crystalline reticulum, brings about a high risk of mechanical destruction of cell structures situated along the path of crystal growth. This occurs both outside the cells, in the solutions where they are placed for the freezing phase, and inside the cells themselves. This drawback is only partially overcome by using cryopreserving agents such as glycerine, dimethylsulfoxide etc.

The very high pressures acting on the cell structures when water passes from the liquid to the solid stage act synergistically with the mechanical damage deriving from the formation of ice needles.

The volume increase for aqueous solutions is up to 10%.

Both the crystal formation and the volume change at the liquid-solid transition phase usually occur in the solutions presently used for the cryopreservation of biological material and are the main cause of the

viability loss of the cryopreserved samples. The present invention provides cryopreserving solutions in which the liquid-solid transition phase occurs without volume increase and macrocrystal formation, providing therefore improvements over the prior-art methods and solving in an economic and convenient way the many problems of the cryopreservation of biological material, particularly those previously cited for the preservation of frozen epithelial tissue sheets.

The solutions of the invention comprise:

- a) polyethylene glycols (PEG) having molecular weight not higher than 20 KD, preferably from 3 to 5 KD, at concentrations not higher than 25% w/w, preferably from 10 to 20% w/w;
- b) at least one low molecular weight compound, defined as a cross-linking agent in view of its ability to form with the polymer and with water cross-linked systems based on hydrogen bonds, in concentrations not higher than 25% w/w, preferably from 5 to 20% w/w. The cross-linking agents are selected from polyols (glycerine, ethylene glycols, maltitol, inositol), mono and oligosaccharides (glucose, sucrose, maltodextrins) and PEGs having molecular weight lower than 1 KD. Glycerine is preferably used.

The solutions of the invention may be prepared in physiological saline solutions and may contain antibiotics, proteins, sera and other compounds usually used for the growth and culture of biological materials.



More particularly, solutions suited for the cryopreservation of viable, in vitro grown epithelium sheets are prepared by adding in suitable concentrations PEG and glycerol as cross-linking agent, preferably 13% w/v PEG and 15% w/v glycerol to media optimized for the keratinocytes growth such as 60% v/v Dulbecco modification of Eagle's medium, 30% v/v Ham's F12, 10% v/v Foetal calf serum (FCS), containing glutamine 4 mM, adenine  $1.8 \times 10^{-4}$  M, insuline 5 µg/mL, transferrine 5 µg/mL, triiodothyronina  $2 \times 10^{-9}$  M hydrocortisone 0.4 µg/mL, penicillin-streptomycine 50 U/mL.

The sheets are previously washed in the cryopreserving solution and then introduced into suitable bags containing 100-200 ml of solution.

After closing the containers, the samples are allowed to equilibrate at room temperature for periods not longer than 1 h, preferably from 5 to 10 min.

Freezing of samples may be carried out by simply transferring the containers at the preservation temperature, generally kept from -25 to -100°C, preferably -80°C.

Thawing of samples may be carried out by dipping the frozen container in a bath at 37°C for 5-10 min. In order to make this step less critical, especially when not particularly qualified users have to handle such delicate cellular materials, and to minimize stress to which thawed cells may be subjected by the high concentration of cryoprotective agents in the liquid phase, thawing of epithelial sheets may be carried out

in a special container consisting of two compartments which are interconnected when the sample is thawed, one containing the epithelial sheets in the cryopreserving solution and the other, containing nutrient medium or isotonic saline solution. During thawing, the two solutions are mixed, causing the dilution of cryoprotective agents minimizing thereby possible cellular damage.

The human epithelial sheets cryopreserved in the solutions of the invention are equivalent to the fresh ones not subjected to freezing, both as far as the morphology and the after graft are concerned.

The solutions of the invention may also be used for the preservation of cells used for the preparation of the sheets.

These comprise, for instance, 3T3 cells to be used as feeder layer or keratinocytes isolated and dissociated from biopsies.

The following examples further illustrate the invention.

#### EXAMPLE 1

Keratinocytes cultures were obtained according to the method of Rheinwald and Green. The secondary confluent keratinocytes cultures obtained according to US 4016036 are detached from culture flasks, transferred on suitable support and anchored by means of metal clips for vascular surgery according to the method of US 4304866.

The so prepared sheets are washed by immersion in sterile conditions in 100 ml of the cryopreserving

solution consisting of 60% v/v Dulbecco modification of Eagle's medium, 30% v/v Ham's F12, 10% v/v FCS, glutamine 4 mM, adenine  $1.8 \times 10^{-4}$  M, insuline 5  $\mu$ g/mL, transferrine 5  $\mu$ g/mL, triiodothyrosine  $2 \times 10^{-9}$  M, hydrocortisone 0.4  $\mu$ g/mL, EGF 10 ng/mL, penicilline-streptomycine 50 U/mL, PEG 3350 13% p/v, glycerol 15% p/v.

The sheets are then transferred under sterile conditions in 15 x 10 cm plastic envelopes, previously sterilized and containing 100 ml of the above reported cryopreserving solution.

The plastic envelopes are then heat sealed and then transferred, without any particular precaution, to the freezer and cooled to a temperature ranging from -25 and -100°C, preferably -80°C. The duration of the operations bringing the epithelial sheets into contact with the cryopreserving solution at room temperature is not critical and may range from a few minutes to 1 h, preferably 5-10 min, without negative consequences in the clinical use of the preserved material.

The liquid-solid transition phase of the solution occurs between -10 and -15°C and the cooling rate of the samples is not critical as well as the constance of the preserving temperature which may oscillate even by tens of degrees without affecting viability of the preserved material.

The epithelial sheets may be preserved at -80°C for several months, whereas at higher temperatures this period is slightly shortened. The envelopes containing the frozen epithelial sheets may be shipped in solid

CO<sub>2</sub> or at higher temperatures up to -20°C and the therapeutic use is not impaired by temperature changes of tens of degrees or by a short accidental thawing.

Immediately before use, the envelopes, taken out from the freezer are immediately introduced into a water bath at 37°C for 5-10 min and then immersed for a few seconds in 70% ethanol. After opening in sterile conditions, the epithelial sheets are transferred into sterile containers, thoroughly washed with medium free from serum and additives or with saline solution and then used.

The thawed samples are subjected to the following characterizations:

- a) histology of the sheet, morphological exam and evaluation of the keratins typical of the various differentiative degrees;
- b) trypsinization and low density plating to evaluate the number and size of the formed colonies;
- c) microscopic evaluation of the cellular morphology and of the number of viable cells;
- d) biological tests of graftability on derma or other vital receiving bed, such as nude mice.

The overall results of these characterizations show that cryopreserved epithelial sheets in the solutions of the invention remain viable and show characteristics comparable to those of fresh sheets which have been not cryopreserved.

#### EXAMPLE 2

Epithelial sheets cryopreserved at -80°C for 30 days, as disclosed in Example 1, are thawed by

immersing for 5-7 min the container in a bath at 37°C and, after opening in sterile conditions, the epithelial sheets are washed in 100 ml of culture medium consisting of 60% v/v Dulbecco modification of Eagle's medium, 30% v/v Ham's F12, 10% v/v FCS, glutamine 4 mM, adenine  $1.8 \times 10^{-4}$  M, insuline 5 µg/mL, transferrine 5 µg/mL, triodothyronine  $2 \times 10^{-9}$  M, hydrocortisone 0.4 µg/mL, EGF 10 ng/mL, penicillin-streptomycine 50 U/mL.

10        The sheets are then treated with trypsin and dissociated into single cells which are cultivated on a bed of irradiated fibroblasts, as disclosed in US 4016036. When keratinocytes colonies are subconfluent the culture medium is removed, the cell layer is washed with FCS-free culture medium and then treated with  
15        trypsin to dissociate colonies into single cells.

      The obtained cellular suspension is centrifuged, suspended again in 5 ml of cryoprotective solution, centrifuged again at 1000 g for 5 min, the solution is removed, the cell pellet is suspended in 1.5 ml of  
20        cryoprotective solution and frozen to -80°C in three aliquots. At different time intervals, ranging from 1 week to 3 months the cells are thawed placing the closed vial for 5-7 min in a bath at 37°C, 5 ml of  
25        fresh medium are added, the suspension is centrifuged at 1000 g for 5 min and suspended again in the medium with which the culture on irradiated fibroblasts is started, as reported in US 4016036 until a multi-layer epithelium is obtained on which the following  
30        characterizations are carried out:

- a) histology of the sheet, morphological exam and evaluation of the keratins typical of the various differentiative degrees;
- b) trypsinization and low density plating to evaluate the number and size of the formed colonies;
- c) microscopic evaluation of the cellular morphology and of the number of viable cells;
- d) biological tests of graftability on derma or other vital receiving bed, such as nude mice.

The several results of these characterizations show that the cells deriving from epithelium sheets cryopreserved in the solutions of the invention, after enzymatic treatment and freezing, remain perfectly viable and keep their characteristics and differentiative features, as shown by the ability of yielding again epithelial sheets fully comparable to sheets obtained from non frozen cells.

CLAIMS

1. Cryoprotective aqueous solutions useful for the preservation of in vitro cultured epithelial sheets comprising:
  - a) polyethyleneglycols having molecular weight not higher than 20 KD;
  - b) at least one cross-linking agent selected from polyols, polyamines, mono- or oligosaccharides, polyethyleneglycols having molecular weight lower than 1 KD.
2. Solutions according to claim 1 wherein the cross-linking agent b) is selected from glycerol, maltitol, glucose, fructose, sucrose, maltodextrins.
3. Solutions according to claim 1 or 2 wherein the polyethyleneglycol is present in a concentration not higher than 25% w/v, preferably from 10 to 20%.
4. Solutions according to any one of the previous claims wherein the cross-linking agent is used at a concentration not higher than 25% w/v, preferably from 5 to 25%.
5. Solutions according to any one of the previous claims also containing antibiotics, proteins, sera and other usual components used for the growth and culture of keratinocytes and sheets of cultured epithelium.
6. Solutions according to any one of the previous claims comprising 60% v/v Dulbecco modification of Eagle's medium, 30% v/v Ham's F12, 10% v/v Foetal calf serum (FCS), containing glutamine 4 mM, adenine 1.8 x

$10^{-4}$  M, insuline 5  $\mu\text{g/mL}$ , transferrine 5  $\mu\text{g/mL}$ , triiodothyronina  $2 \times 10^{-9}$  M, hydrocortisone 0,4  $\mu\text{g/mL}$ , Epidermal Growth Factor (EGF) 10  $\text{ng/mL}$ , penicillin-streptomycine 50 U/mL PEG and a cross-linking agent.

7. A method for the cryopreservation of epithelial sheets obtained from keratinocytes cultures comprising:

- a) incubation at room temperature of said epithelial sheets in a cryopreserving solution of claims 1-6, for a time shorter than 1 h, preferably 5-15 min;
- b) freezing in said solution at temperatures from  $-100$  to  $-25^{\circ}\text{C}$ .

8. A method for the preservation of cells used for the preparation of epithelial sheets expanded in vitro such as 3T3 cells or isolated keratinocytes which comprising:

- a) incubation at room temperature of said cells in a cryopreserving solution of claims 1-6, for a time shorter than 1 h, preferably 5-15 min;
- b) freezing in said solution at temperatures from  $-100$  to  $-25^{\circ}\text{C}$ .

9. Plastic envelopes containing frozen epithelial sheets obtained according to the method of claim 7.

10. Epithelial sheets obtainable by the process of claim 7.



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 93/03364

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 5 A01N1/02 A01N25/34

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 5 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,91 18505 (BIOSURFACE TECHNOLOGY INC) 12 December 1991 see page 7, line 1 - page 8, line 18; claims; examples ---	1-9
X	TRANSPLANTATION vol. 49, no. 2, February 1990, USA pages 261 - 264 W.N.WICOMB ET. AL. 'Optimal cardioplegia and 24-hour heart storage with simplified UW solution containing polyethylene glycol' see Materials and Methods, Table 1 --- -/--	1,3,4

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "A" document member of the same patent family

Date of the actual completion of the international search

9 March 1994

Date of mailing of the international search report

16.03.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Donovan, T

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 93/03364

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,93 00807 (CRYOLIFE INC) 21 January 1993 see page 3, line 34 - page 3, line 13 see page 6, line 10 - page 8, line 5; claims; figures; examples ---	1-9
P,X	WO,A,93 14191 (CRYOPHARM CORPORATION) 22 July 1993 see page 3, line 17 - page 9, line 17 see page 17, line 26 - page 20, line 21; claims; examples ---	1-9
A	CRYO-LETTERS vol. 3, 1982 pages 118 - 120 F.FRANKS 'apparent osmotic activities of water soluble polymers used as cryoprotectants' ---	
A	WO,A,91 11101 (OSTEOTECH, INC) 8 August 1991 ---	
A	WO,A,92 20300 (CRYOLIFE, INC) 26 November 1992 ---	
A	US,A,4 086 218 (E.SHANBROM ET. AL.) 25 April 1978 ---	
A	US,A,4 105 650 (E.SHANBROM ET. AL.) 8 August 1978 ---	
A	EP,A,0 296 475 (ISTITUTO NAZIONALE PER LA RICERCA SUL CANCRO) 28 December 1988 cited in the application ---	
A	EP,A,0 364 306 (TUTORTEC INC.) 18 April 1990 -----	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 93/03364

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9118505	12-12-91	US-A- 5145770 AU-A- 8060191 EP-A- 0532670	08-09-92 31-12-91 24-03-93
WO-A-9300807	21-01-93	AU-A- 2309692	11-02-93
WO-A-9314191	22-07-93	AU-B- 3443093	03-08-93
WO-A-9111101	08-08-91	US-A- 5118512	02-06-92
WO-A-9220300	26-11-92	US-A- 5160313 AU-A- 1891892 CA-A- 2103025 EP-A- 0584195	03-11-92 30-12-92 15-11-92 02-03-94
US-A-4086218	25-04-78	US-A- 4105650	08-08-78
US-A-4105650	08-08-78	US-A- 4086218	25-04-78
EP-A-0296475	28-12-88	AU-A- 1995788 DE-A- 3874019 WO-A- 8810068 JP-T- 2500800 OA-A- 9038	19-01-89 01-10-92 29-12-88 22-03-90 31-03-91
EP-A-0364306	18-04-90	AU-A- 4277689 CA-A- 2000181 JP-A- 2174848	26-04-90 14-04-90 06-07-90

**This Page Blank (uspio)**